

The Indefinite Self-Dual Metrics and Painleve Equations

Shoji Okumura (okumura@math.sci.osaka-u.ac.jp)
Math. Sci. Osaka Univ., Japan

Abstract

We classify the $SU(2)$ -invariant anti-self-dual metrics with a signature $(+, +, -, -)$. The metrics are specified by a solution of Painlevé VI, V, III or II. Moreover we show the geometric meaning of the metrics specified by each type of Painlevé functions.

1 Introduction

The aim of this paper is to classify the anti-self-dual metrics in real dimension four admitting an isometric action of $SU(2)$ with generically three-dimensional orbits. In this paper, we study not only the definite metrics, but also the indefinite metrics with a signature $(+, +, -, -)$.


Hitchin [7] shows that the $SU(2)$ -invariant anti-self-dual metric is generically specified by a solution of Painlevé VI with two complex parameters. He used the twistor correspondence [10] to associate the anti-self-dual equation and Painlevé equation. On twistor space, the lifted action of $SU(2)$ determines a pre-homogenous action of $SU(2)$, and it determines a isomonodromic family of connections on $\mathbb{CP}^1$, and then we have Painlevé equations. In this framework, Dancer [4] shows that the diagonal scalar-flat Kähler metric is specified by a solution of Painlevé III with parameters $(4, 4, 4 - 4)$. The author [8] shows that the $SU(2)$ -invariant anti-self-dual hermitian metric is specified by a solution of Painlevé III with one complex parameter.

If the metric is definite, then the anti-self-dual equation reduce to either Painlevé VI or III. On the other hand, we show that, if the metric has a signature $(+, +, -, -)$, then the anti-self-dual equation reduce to not only Painlevé VI or III but also Painlevé V or II. The difference of types of Painlevé is due to the difference of reality on the twistor space.

Painlevé VI is shown to be deformation equations for a linear problem

$$\left(\frac{d}{dz} - B_1 \right) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = 0,$$

where B_1 has four simple poles on $\mathbb{CP}^1$ [5]. And Painleve II, III, IV, V are degenerated from Painlevé VI:



This is the confluence diagram of poles of B_1 , where the Roman numerals represent the types of Painlevé, and the parenthesized numbers represent the orders of poles of B_1 . For example, Painlevé V is shown to be deformation equations for a linear problem with one double and two simple poles. Due to the reality of twistor space, the poles of B_1 makes two conjugate pairs. Therefore, the configuration of poles never becomes Painlevé IV type.

The multiple pole of B_1 determines a hermitian structure or a structure of null surfaces, and this is the geometric meaning of the metrics specified by each type of Painlevé functions.

Acknowledgements. The author would express his sincere gratitude to Professor Yousuke Ohyama, who introduced him to the subject, for enlightening discussions.

2 The Diagonal Anti-self-dual Equations

In this section, we review the anti-self-dual equations on the $SU(2)$ -invariant diagonal metrics.

The $SU(2)$ -invariant diagonal metric is written in the following form:

$$g = w_1 w_2 w_3 dt^2 + \frac{w_2 w_3}{w_1} \sigma_1^2 + \frac{w_3 w_1}{w_2} \sigma_2^2 + \frac{w_1 w_2}{w_3} \sigma_3^2. \quad (1)$$

w_1, w_2 and w_3 are functions of t , and $\sigma_1, \sigma_2, \sigma_3$ are left invariant one-forms on each $SU(2)$ -orbit satisfying

$$d\sigma_1 = \sigma_2 \wedge \sigma_3, \quad d\sigma_2 = \sigma_3 \wedge \sigma_1, \quad d\sigma_3 = \sigma_1 \wedge \sigma_2. \quad (2)$$

Tod [11] showed that the anti-self-dual equations on the $SU(2)$ -invariant diagonal metric are given by the following system:

$$\begin{aligned} \dot{w}_1 &= -w_2 w_3 + w_1 (\alpha_2 + \alpha_3), \\ \dot{w}_2 &= -w_3 w_1 + w_2 (\alpha_3 + \alpha_1), \\ \dot{w}_3 &= -w_1 w_2 + w_3 (\alpha_1 + \alpha_2), \\ \dot{\alpha}_1 &= -\alpha_2 \alpha_3 + \alpha_1 (\alpha_2 + \alpha_3), \\ \dot{\alpha}_2 &= -\alpha_3 \alpha_1 + \alpha_2 (\alpha_3 + \alpha_1), \\ \dot{\alpha}_3 &= -\alpha_1 \alpha_2 + \alpha_3 (\alpha_1 + \alpha_2), \end{aligned} \quad (3)$$

where $\alpha_1, \alpha_2, \alpha_3$ are auxiliary functions and the dots denote differentiation with respect to t . The anti-self-dual equation (3) has a first integral

$$k = \frac{\alpha_1(w_2^2 - w_3^2) + \alpha_2(w_3^2 - w_1^2) + \alpha_3(w_1^2 - w_2^2)}{8(\alpha_1 - \alpha_2)(\alpha_2 - \alpha_3)(\alpha_3 - \alpha_1)}.$$

Furthermore, if we set

$$x = \frac{\alpha_2 - \alpha_1}{\alpha_2 - \alpha_3},$$

$$q = \frac{w_2(\alpha_1 - \alpha_2)(w_2(w_1^2 - w_3^2) + 2\sqrt{2k}w_1w_3(\alpha_1 - \alpha_3))}{w_1^2(w_2^2 - w_3^2)\alpha_1 + w_2^2(w_3^2 - w_1^2)\alpha_2 + w_3^2(w_1^2 - w_2^2)\alpha_3},$$

then the system (3) generically reduces to a family of Painlevé VI with special parameters

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{(\sqrt{2k} - 1)^2}{2}, k, -k, \frac{1 + 2k}{2} \right).$$

We will review the Painlevé equations in the appendix.

3 The Non-diagonal Anti-self-dual Equations

We can write a $SU(2)$ -invariant metric in the form

$$g = f(\tau)d\tau^2 + \sum_{l,m=1}^3 h_{lm}(\tau) \sigma_l \sigma_m.$$

Using the Killing form, we can diagonalize the metric g on each $SU(2)$ -orbit. Then we can express the metric as follows:

$$g = (abc)^2 dt^2 + a^2 d\hat{\sigma}_1^2 + b^2 d\hat{\sigma}_2^2 + c^2 d\hat{\sigma}_3^2,$$

for some $t = t(\tau)$, $a = a(t)$, $b = b(t)$, $c = c(t)$ and

$$\begin{pmatrix} \hat{\sigma}_1 \\ \hat{\sigma}_2 \\ \hat{\sigma}_3 \end{pmatrix} = R(t) \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix},$$

where $R(t)$ is $SO(3)$ -valued function.

Since $\dot{R}R^{-1} \in \mathfrak{so}(3)$, we can write

$$d \begin{pmatrix} \hat{\sigma}_1 \\ \hat{\sigma}_2 \\ \hat{\sigma}_3 \end{pmatrix} = R(t) \begin{pmatrix} \sigma_2 \wedge \sigma_3 \\ \sigma_3 \wedge \sigma_1 \\ \sigma_2 \wedge \sigma_2 \end{pmatrix} + \dot{R} dt \wedge \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix}$$

$$= \begin{pmatrix} \hat{\sigma}_2 \wedge \hat{\sigma}_3 \\ \hat{\sigma}_3 \wedge \hat{\sigma}_1 \\ \hat{\sigma}_1 \wedge \hat{\sigma}_2 \end{pmatrix} + \begin{pmatrix} 0 & \xi_3 & -\xi_2 \\ -\xi_3 & 0 & \xi_1 \\ \xi_2 & -\xi_1 & 0 \end{pmatrix} dt \wedge \begin{pmatrix} \hat{\sigma}_1 \\ \hat{\sigma}_2 \\ \hat{\sigma}_3 \end{pmatrix},$$

for some $\xi_1 = \xi_1(t)$, $\xi_2 = \xi_2(t)$, $\xi_3 = \xi_3(t)$.

If $\xi_1 = 0$, $\xi_2 = 0$, $\xi_3 = 0$, then the matrix (h_{lm}) can be chosen to be diagonal for all τ , and then we say that g is a diagonal metric.

In the following, we mainly study the non-diagonal case.

We set $w_1 = bc$, $w_2 = ca$, $w_3 = ab$ and determine $\alpha_1, \alpha_2, \alpha_3$ by

$$\begin{aligned} \dot{w}_1 &= -w_2w_3 + w_1(\alpha_2 + \alpha_3), \\ \dot{w}_2 &= -w_3w_1 + w_2(\alpha_3 + \alpha_1), \\ \dot{w}_3 &= -w_1w_2 + w_3(\alpha_1 + \alpha_2). \end{aligned} \tag{4}$$

Then the anti-self-dual equations are as follows [8]:

$$\begin{aligned}
\dot{\alpha}_1 &= -\alpha_2\alpha_3 + \alpha_1(\alpha_2 + \alpha_3) + \frac{1}{4}(w_2^2 - w_3^2)^2 \left(\frac{\xi_1}{w_2w_3} \right)^2 \\
&\quad + \frac{1}{4}(w_3^2 - w_1^2)(3w_1^2 + w_3^2) \left(\frac{\xi_2}{w_3w_1} \right)^2 \\
&\quad + \frac{1}{4}(w_2^2 - w_1^2)(3w_1^2 + w_2^2) \left(\frac{\xi_3}{w_1w_2} \right)^2, \\
\dot{\alpha}_2 &= -\alpha_3\alpha_1 + \alpha_2(\alpha_3 + \alpha_1) + \frac{1}{4}(w_3^2 - w_1^2)^2 \left(\frac{\xi_2}{w_3w_1} \right)^2 \\
&\quad + \frac{1}{4}(w_1^2 - w_2^2)(3w_2^2 + w_1^2) \left(\frac{\xi_3}{w_1w_2} \right)^2 \\
&\quad + \frac{1}{4}(w_3^2 - w_2^2)(3w_2^2 + w_3^2) \left(\frac{\xi_1}{w_2w_3} \right)^2, \\
\dot{\alpha}_3 &= -\alpha_1\alpha_2 + \alpha_3(\alpha_1 + \alpha_2) + \frac{1}{4}(w_1^2 - w_2^2)^2 \left(\frac{\xi_3}{w_1w_2} \right)^2 \\
&\quad + \frac{1}{4}(w_2^2 - w_3^2)(3w_3^2 + w_2^2) \left(\frac{\xi_1}{w_2w_3} \right)^2 \\
&\quad + \frac{1}{4}(w_1^2 - w_3^2)(3w_3^2 + w_1^2) \left(\frac{\xi_2}{w_3w_1} \right)^2,
\end{aligned} \tag{5}$$

and

$$\begin{aligned}
(w_2^2 - w_3^2) \frac{d}{dt} \left(\frac{\xi_1}{w_2w_3} \right) &= \frac{\xi_2}{w_3w_1} \frac{\xi_3}{w_1w_2} (-2w_2^2w_3^2 + w_3^2w_1^2 + w_1^2w_2^2) \\
&\quad + \frac{\xi_1}{w_2w_3} (\alpha_2w_2^2 - \alpha_3w_3^2 + 3\alpha_2w_3^2 + 3\alpha_3w_2^2), \\
(w_3^2 - w_1^2) \frac{d}{dt} \left(\frac{\xi_2}{w_3w_1} \right) &= \frac{\xi_3}{w_1w_2} \frac{\xi_1}{w_2w_3} (-2w_3^2w_1^2 + w_1^2w_2^2 + w_2^2w_3^2) \\
&\quad + \frac{\xi_2}{w_3w_1} (\alpha_3w_3^2 - \alpha_1w_1^2 + 3\alpha_3w_1^2 + 3\alpha_1w_2^2), \\
(w_1^2 - w_2^2) \frac{d}{dt} \left(\frac{\xi_3}{w_1w_2} \right) &= \frac{\xi_1}{w_1w_3} \frac{\xi_2}{w_3w_1} (-2w_1^2w_2^2 + w_2^2w_3^2 + w_3^2w_1^2) \\
&\quad + \frac{\xi_3}{w_1w_2} (\alpha_1w_1^2 - \alpha_2w_2^2 + 3\alpha_1w_2^2 + 3\alpha_2w_1^2).
\end{aligned} \tag{6}$$

Remark 3.1 If $\xi_1 = 0$, $\xi_2 = 0$ and $\xi_3 = 0$ then the system of equations (4), (5) and (6) reduces to a sixth-order system (3) given by Tod [11]. Furthermore, if $\alpha_1 = w_1, \alpha_2 = w_2, \alpha_3 = w_3$ then (4),(5),(6) reduce to a third-order system which determines Atiyah-Hitchin family [1], and if $\alpha_1 = 0, \alpha_2 = 0, \alpha_3 = 0$ then the system reduces to a third-order system which determines BGPP family [3].

Remark 3.2 If $w_2 = w_3$, then we can set $\xi_1 = 0$, $\xi_2 = 0$ and $\xi_3 = 0$ by taking another frame. This is also a diagonal case. Therefore we assume $(w_2 - w_3)(w_3 - w_1)(w_1 - w_2) \neq 0$.

4 The Isomonodromic Deformations

Let (M, g) be an oriented Riemannian four manifold. We define a manifold Z to be the unit sphere bundle in the bundle of self-dual two-forms, and let $\pi : Z \rightarrow M$ denote the projection. Each point z in the fiber over $\pi(z)$ defines a complex structure on the tangent space $T_{\pi(z)}M$, compatible with the metric and its orientation.

Using the Levi-Civita connection, we can split the tangent space $T_z Z$ into horizontal and vertical spaces, and the projection π identifies the horizontal space with $T_{\pi(z)}M$. This space has a complex structure defined by z and the vertical space is the tangent space of the fiber $S^2 \cong \mathbb{CP}^1$ which has its natural complex structure.

The almost complex structure on Z is integrable if and only if the metric is anti-self-dual [2, 10]. In this situation Z is called the twistor space of (M, g) and the fibers are called the real twistor lines.

The almost complex structure on Z can be determined by the following $(1, 0)$ -forms:

$$\begin{aligned}\Theta_1 &= z(e^1 + \sqrt{-1}e^2) - (e^0 + \sqrt{-1}e^3), \\ \Theta_2 &= z(e^0 - \sqrt{-1}e^3) + (e^1 - \sqrt{-1}e^2), \\ \Theta_3 &= dz + \frac{1}{2}z^2(\omega_3^0 - \omega_2^1 + \sqrt{-1}(\omega_1^0 - \omega_3^2)) \\ &\quad - \sqrt{-1}z(\omega_2^0 - \omega_1^3) + \frac{1}{2}(\omega_1^0 - \omega_2^1 - \sqrt{-1}(\omega_1^0 - \omega_3^2)),\end{aligned}\tag{7}$$

where $\{e^0, e^1, e^2, e^3\}$ is an orthonormal frame, and ω_j^i are the connection forms determined by $de^i + \omega_j^i \wedge e^j = 0$ and $\omega_j^i + \omega_i^j = 0$. Then the anti-self-dual condition is

$$d\Theta_1 \equiv 0, \quad d\Theta_2 \equiv 0, \quad d\Theta_3 \equiv 0 \quad (\text{mod } \Theta_1, \Theta_2, \Theta_3).\tag{8}$$

Theorem 4.1 *If the metric is positive definite, then the Pfaffian is invariant under conjugate action and $z \rightarrow -1/\bar{z}$ [2]. If the metric has a signature $(+, +, -, -)$, then the Pfaffian is invariant under conjugate action and $z \rightarrow \bar{z}$.*

If the metric is $SU(2)$ invariant, we can write

$$\begin{pmatrix} \Theta_1 \\ \Theta_2 \\ \Theta_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} dz + \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} dt + A \begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix},\tag{9}$$

where $v_1 = v_1(z, t)$, $v_2 = v_2(z, t)$, $v_3 = v_3(z, t)$; $A = (a_{ij}(z, t))_{i,j=1,2,3}$.

If $\det A \equiv 0$, then metric turns to be diagonal, and the metric is in the BGPP family [3].

If $\det A \neq 0$, then we can write

$$\begin{pmatrix} \sigma_1 \\ \sigma_2 \\ \sigma_3 \end{pmatrix} \equiv -A^{-1} \left(\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} dz + \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} dt \right), \quad \text{mod } \Theta_1, \Theta_2, \Theta_3.\tag{10}$$

If we set

$$\begin{pmatrix} s_1 \\ s_2 \\ s_3 \end{pmatrix} := -A^{-1} \left(\begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} dz + \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} dt \right), \quad (11)$$

then

$$d \begin{pmatrix} s_1 \\ s_2 \\ s_3 \end{pmatrix} \equiv \begin{pmatrix} s_2 \wedge s_3 \\ s_3 \wedge s_1 \\ s_1 \wedge s_2 \end{pmatrix}, \quad \text{mod } \Theta_1, \Theta_2, \Theta_3. \quad (12)$$

Since s_1, s_2, s_3 are one-forms on (z, t) -plane, the congruency equation (12) turns to be a plain equation:

$$d \begin{pmatrix} s_1 \\ s_2 \\ s_3 \end{pmatrix} = \begin{pmatrix} s_2 \wedge s_3 \\ s_3 \wedge s_1 \\ s_1 \wedge s_2 \end{pmatrix}. \quad (13)$$

If the metric is positive definite, then s_1, s_2, s_3 are invariant under conjugate action and $z \rightarrow -1/\bar{z}$ by theorem 4.1. And if the metric has a signature $(+, +, -, -)$, then s_1, s_2, s_3 are invariant under conjugate action and $z \rightarrow \bar{z}$.

If we set

$$\Sigma = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{-1}s_1 & -s_3 + \sqrt{-1}s_2 \\ s_3 + \sqrt{-1}s_2 & -\sqrt{-1}s_1 \end{pmatrix} \quad (14)$$

$$=: -B_1 dz - B_2 dt, \quad (15)$$

then

$$d\Sigma + \Sigma \wedge \Sigma = 0. \quad (16)$$

This is the isomonodromy condition for the the following linear problem [5]

$$\left(\frac{d}{dz} - B_1 \right) \begin{pmatrix} y_1 \\ y_2 \end{pmatrix} = 0. \quad (17)$$

Lemma 4.2 *The components of B_1 are rational functions of z ,*

$$B_1 = \frac{F(z)}{G(z)},$$

where $F(z)$ is degree 2 and $G(z)$ is degree 4. If the metric is positive definite, then $B_1 \rightarrow -{}^t B_1$ under conjugate action and $z \rightarrow -1/\bar{z}$. And if the metric has a signature $(+, +, -, -)$, then $B_1 \rightarrow -{}^t B_1$ under conjugate action and $z \rightarrow \bar{z}$.

For this lemma, generically B_1 has four simple poles. In this case, the deformation equation of (17) is Painlevé VI.

Theorem 4.3 *The anti-self-dual equations on $SU(2)$ -invariant metrics generically reduce to Painlevé VI.*

The idea of Hitchin [7] is that the lifted action of $SU(2)$ on the twistor space Z gives a homomorphism of vector bundles $\alpha : Z \times \mathfrak{su}(2)^\mathbb{C} \rightarrow TZ$, and the inverse of α gives a flat meromorphic $SL(2, \mathbb{C})$ -connection, which determine isomonodromic deformations. Since one-forms $\Theta_1, \Theta_2, \Theta_3$ on Z can be considered as infinitesimal variations, we can identify Σ with α^{-1} .

First, we review the positive definite metric. By lemma 4.2 the poles of B_1 make antipodal pairs $\zeta_0, -1/\bar{\zeta}_0$, and $\zeta_1, -1/\bar{\zeta}_1$ on $\mathbb{CP}^1$.

Therefore, we have two types of configuration of poles of B_1 :

- (a) B_1 has four simple poles $\zeta_0, \bar{\zeta}_0, \zeta_1, \bar{\zeta}_1$ on $\mathbb{C} \setminus \mathbb{R}$.

$$B_1 = \frac{A_0}{z - \zeta_0} + \frac{-{}^t\bar{A}_0}{z + 1/\bar{\zeta}_0} + \frac{A_1}{z - \zeta_1} + \frac{-{}^t\bar{A}_1}{z + 1/\bar{\zeta}_1}.$$

The deformation equation is Painlevé VI with parameters,

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{1}{2}(\theta_0 - 1)^2, \frac{1}{2}\bar{\theta}_0^2, -\frac{1}{2}\theta_1^2, \frac{1}{2}(1 + \bar{\theta}_1^2) \right),$$

where $\theta_0^2 = 2 \operatorname{tr} A_0^2$, $\theta_1^2 = 2 \operatorname{tr} A_1^2$.

- (b) B_1 has two double poles $\zeta, \bar{\zeta}$ on $\mathbb{C} \setminus \mathbb{R}$.

$$B_1 = \frac{A_2}{(z - \zeta)^2} + \frac{\sqrt{-1}C}{z - \zeta} + \frac{-\sqrt{-1}C}{z + 1/\bar{\zeta}} + \frac{-{}^t\bar{A}_2/\bar{\zeta}^2}{(z + 1/\bar{\zeta})^2},$$

where $C = -{}^t\bar{C}$. The deformation equation is Painlevé III with parameters,

$$(\alpha, \beta, \gamma, \delta) = (4\theta, 4(1 + \bar{\theta}), 4, -4),$$

where $\theta^2 = 2(\operatorname{tr}(A_3 C))^2 / \operatorname{tr} C^2$.

Theorem 4.4 *If the metric is positive definite, the anti-self-dual equations reduce to the following two Painlevé equations:*

- (a) *A family of Painlevé VI with two complex parameters,*

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{1}{2}(\theta_0 - 1)^2, \frac{1}{2}\bar{\theta}_0^2, -\frac{1}{2}\theta_1^2, \frac{1}{2}(1 + \bar{\theta}_1^2) \right),$$

- (b) *A family of Painlevé III with one complex parameter,*

$$(\alpha, \beta, \gamma, \delta) = (4\theta, 4(1 + \bar{\theta}), 4, -4).$$

Remark 4.5 *Hitchin shows the anti-self-dual equations reduce to Painlevé VI with the parameters above ([7], P.50 (14)). Dancer [4] shows the scalar-flat diagonal Kähler metric is specified by a solution of Painlevé III with parameters $(\alpha, \beta, \gamma, \delta) = (4, 4, 4, -4)$. Now, theorem 4.4 (b) is a generalization of Dancer's result.*

From now on, we will classify the anti-self-dual metrics with a signature $(+, +, -, -)$. By lemma 4.2 the poles of B_1 make conjugate pairs $\zeta_0, \bar{\zeta}_0$, and $\zeta_1, \bar{\zeta}_1$ in $\mathbb{CP}^1$. Therefore we have five types of metrics corresponding to configuration of poles of B_1 .

Theorem 4.6 *If the metric has a signature $(+, +, -, -)$, the anti-self-dual equations reduce to the following five Painlevé equations:*

(a) *Painlevé VI with two complex parameters*

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{1}{2}(\theta_0 - 1)^2, \frac{1}{2}\bar{\theta}_0^2, -\frac{1}{2}\theta_1^2, \frac{1}{2}(1 + \bar{\theta}_1^2) \right),$$

(b) *Painlevé V with one real and one complex parameters*

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{1}{2}(\theta_0 + \bar{\theta}_0 + \theta_\infty)^2, -\frac{1}{2}(\theta_0 + \bar{\theta}_0 - \theta_\infty)^2, 1 - \theta_0 + \bar{\theta}_0, \frac{1}{2} \right),$$

where $\theta_\infty \in \mathbb{R}$.

(c) *Painlevé III with one complex parameter*

$$(\alpha, \beta, \gamma, \delta) = (4\theta, 4(1 + \bar{\theta}), 4, -4).$$

(d) *Painlevé III with two real parameters*

$$(\alpha, \beta, \gamma, \delta) = (4\theta_1, 4(1 + \theta_2), 4, -4).$$

(e) *Painlevé II with one real parameter α .*

Proof.

Since the poles of B_1 make conjugate pairs $\zeta_0, \bar{\zeta}_0$ and $\zeta_1, \bar{\zeta}_1$, we have five types of configuration of poles of B_1 . In each case, we can calculate local exponents at singularities. These local exponents corresponding to parameters of Painlevé equations (see [6]).

(a) Generically, B_1 has four simple poles $\zeta_0, \bar{\zeta}_0, \zeta_1, \bar{\zeta}_1$ on $\mathbb{C} \setminus \mathbb{R}$.

$$B_1 = \frac{A_0}{z - \zeta_0} + \frac{-{}^t\bar{A}_0}{z - \bar{\zeta}_0} + \frac{A_1}{z - \zeta_1} + \frac{-{}^t\bar{A}_1}{z - \bar{\zeta}_1}.$$

The deformation equations are Painlevé VI with parameters,

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{1}{2}(\theta_0 - 1)^2, \frac{1}{2}\bar{\theta}_0^2, -\frac{1}{2}\theta_1^2, \frac{1}{2}(1 + \bar{\theta}_1^2) \right),$$

where $\theta_0^2 = 2 \operatorname{tr} A_0^2$, $\theta_1^2 = 2 \operatorname{tr} A_1^2$.

(b) If $\zeta_0 = \bar{\zeta}_0 (= \eta)$, then B_1 has two simple poles $\zeta_1, \bar{\zeta}_1$ on $\mathbb{C} \setminus \mathbb{R}$, and one double pole η on $\mathbb{R}$.

$$B_1 = \frac{C}{(z - \eta)^2} + \frac{-A_2 + {}^t\bar{A}_2}{z - \eta} + \frac{A_2}{z - \zeta_1} + \frac{-{}^t\bar{A}_2}{z - \bar{\zeta}_1},$$

where $C = -{}^t\bar{C}$. The deformation equation is Painlevé V with parameters,

$$(\alpha, \beta, \gamma, \delta) = \left(\frac{1}{2}(\theta_0 + \bar{\theta}_0 + \theta_\infty)^2, -\frac{1}{2}(\theta_0 + \bar{\theta}_0 - \theta_\infty)^2, 1 - \theta_0 + \bar{\theta}_0, \frac{1}{2} \right),$$

where $\theta_0^2 = 2 \operatorname{tr} A_2^2$, $\theta_\infty^2 = 2 (\operatorname{tr} (A_2 - {}^t\bar{A}_2) C)^2 / \operatorname{tr} C^2$.

(c) If $\zeta_0 = \zeta_1 (= \zeta)$, then B_1 has two double poles $\zeta, \bar{\zeta}$ on $\mathbb{C} \setminus \mathbb{R}$.

$$B_1 = \frac{A_3}{(z - \zeta)^2} + \frac{\sqrt{-1}C}{z - \zeta} + \frac{-\sqrt{-1}C}{z - \bar{\zeta}} + \frac{-{}^t\bar{A}_3}{(z - \bar{\zeta})^2},$$

where $C = -{}^t\bar{C}$. The deformation equation is Painlevé III with parameters,

$$(\alpha, \beta, \gamma, \delta) = (4\theta, 4(1 + \bar{\theta}), 4, -4),$$

where $\theta^2 = 2(\text{tr}A_3C)^2/\text{tr}C^2$.

(d) If $\zeta_0 = \bar{\zeta}_0 (= \eta_0)$, $\zeta_1 = \bar{\zeta}_1 (= \eta_1)$, then B_1 has two double poles η_0, η_1 on $\mathbb{R}$.

$$B_1 = \frac{C_1}{(z - \eta_0)^2} + \frac{C_2}{z - \eta_0} + \frac{-C_2}{z - \eta_1} + \frac{C_3}{(z - \eta_1)^2},$$

where $C_i = -{}^t\bar{C}_i$ ($i = 1, 2, 3$). The deformation equation is Painlevé III with parameters,

$$(\alpha, \beta, \gamma, \delta) = (4\theta_1, 4(1 + \theta_2), 4, -4),$$

where $\theta_1^2 = 2(\text{tr}C_1C_2)^2/\text{tr}C_2^2$, $\theta_2^2 = 2(\text{tr}C_2C_3)^2/\text{tr}C_2^2$.

(e) If $\zeta_0 = \bar{\zeta}_0 = \zeta_1 = \bar{\zeta}_1 (= \eta)$, then B_1 has one quadruple pole η on $\mathbb{R}$.

$$B_1 = \frac{C_1}{(z - \eta)^4} + \frac{C_2}{(z - \eta)^3} + \frac{C_3}{(z - \eta)^2},$$

where $C_i = -{}^t\bar{C}_i$ ($i = 1, 2, 3$). If $\det C_1 \neq 0$, then the deformation equation is Painlevé II with a parameter,

$$\alpha = \frac{1}{2}(1 + \text{tr}C_2C_3).$$

If $\det C_1 = 0$, then the deformation equation is Painlevé I, but since $C_1 = -{}^t\bar{C}_1$, this never occurs. $\blacksquare$

5 Hermitian Structure and Null Surfaces

In this section we will consider the geometric meaning of the metrics corresponding with each type of Painlevé functions.

Lemma 5.1 *Let g be a anti-self-dual metric. If B_1 has a multiple pole, then the Pfaffian $\Theta_1|_{z=\zeta(t)}$, $\Theta_2|_{z=\zeta(t)}$ is integrable. Conversely, if the Pfaffian $\Theta_1|_{z=\zeta(t)}$, $\Theta_2|_{z=\zeta(t)}$ is integrable for some $z = \zeta(t)$, then B_1 has a multiple pole on $z = \zeta(t)$.*

Proof.

If $z = \zeta(t)$ is a multiple zero of $G(z)$, then $G|_{z=\zeta(t)}$ and $dG|_{z=\zeta(t)}$ must be vanish. Furthermore, $\Theta_3 \equiv 0 \pmod{\Theta_1, \Theta_2, G, dG}$. Therefore, the Pfaffian $\Theta_1|_{z=\zeta(t)}, \Theta_2|_{z=\zeta(t)}$ is integrable.

Conversely, if the Pfaffian $\Theta_1|_{z=\zeta(t)}, \Theta_2|_{z=\zeta(t)}$ is integrable for some $z = \zeta(t)$, then $\Theta_3|_{z=\zeta(t)} \equiv 0 \pmod{\Theta_1, \Theta_2}$. Therefore the denominator G of B_1 has zero on $z = \zeta(t)$. Furthermore, $\Theta_3|_{z=\zeta(t)} \equiv 0 \pmod{\Theta_1, \Theta_2, G}$ is equivalent to $dG|_{z=\zeta(t)} = 0$, therefore $z = \zeta(t)$ is a double pole of B_1 . $\blacksquare$

In [8] we show that if a positive definite $SU(2)$ invariant anti-self-dual metric is corresponding with Painlevé III with one complex parameter $(4\theta, 4(1 + \bar{\theta}), 4, -4)$, then the Pfaffian $\Theta_1|_{z=\zeta(t)}, \Theta_2|_{z=\zeta(t)}$ ($z = \zeta(t)$ is a double pole of B_1) determines a $SU(2)$ invariant hermitian structure.

In the same way, we have the following theorem for the metric with a signature $(+, +, -, -)$.

Theorem 5.2 *If the anti-self-dual equations reduce to Painlevé III with one complex parameter $(4\theta, 4(1 + \bar{\theta}), 4, -4)$, then there exists a $SU(2)$ -invariant hermitian structure. Conversely, if there exists a $SU(2)$ -invariant hermitian structure, then the anti-self-dual equations reduce to Painlevé III with parameters above.*

Definition 5.3 *If $g(X, X) = 0$, then $X \in TM$ is said to be a null direction.*

The Pfaffian $\Theta_1|_{z=\eta(t)}, \Theta_2|_{z=\eta(t)}$ ($\eta(t) \in \mathbb{R}$) determines two dimensional null directions on TM .

Definition 5.4 *Let N be a two dimensional subspace of M . If $g(X, X) = 0$ for any $X \in TN$, then N is called a null surface.*

From lemma 5.1, if $z = \eta(t) \in \mathbb{R}$ is a multiple pole of B_1 , then the Pfaffian $\Theta_1|_{z=\eta(t)}, \Theta_2|_{z=\eta(t)}$ is integrable, and then for any $x \in M$ there exists a $SU(2)$ -invariant null surface passing through x . Conversely, for any $x \in M$, if there exists a $SU(2)$ -invariant null surface passing through x , then the null surface is represented by Pfaffian $\Theta_1|_{z=\eta(t)}, \Theta_2|_{z=\eta(t)}$ for some $\eta(t) \in \mathbb{R}$, and then $z = \eta(t)$ is a multiple pole of B_1 .

Theorem 5.5 *If the anti-self-dual equations reduce to Painlevé V with one real and one complex parameters $(\frac{1}{2}(\theta_0 + \bar{\theta}_0 + \theta_\infty)^2, -\frac{1}{2}(\theta_0 + \bar{\theta}_0 - \theta_\infty)^2, 1 - \theta_0 + \bar{\theta}_0, \frac{1}{2})$, where $\theta_\infty \in \mathbb{R}$, or Painlevé II with one real parameter α , then for any $x \in M$ there exists one $SU(2)$ -invariant null surface passing through x . Conversely, for any $x \in M$ there exists one $SU(2)$ -invariant null surface passing through x , then the anti-self-dual reduce to Painlevé V or II with parameters above.*

If the anti-self-dual equations reduce to Painlevé III with two real parameters $(4\theta_1, 4(1 - \theta_2), 4, -4)$, then for any $x \in M$ there exist two $SU(2)$ -invariant null surfaces passing through x . Conversely, for any $x \in M$ there exist two $SU(2)$ -invariant null surfaces passing through x , then the anti-self-dual reduce to Painlevé III with parameters above.

6 Summary

We classified the $SU(2)$ -invariant anti-self-dual metric with a signature $(+, +, -, -)$ into the five cases (a)–(e) (theorem 4.6). The meaning of the types of Painlevé equations are as follows:

1. Generically, the anti-self-dual metric are specified by a solution of Painlevé VI with two complex parameters.

2. If the anti-self-dual metric is specified by a solution of Painlevé III with two complex parameters, then there exist a $SU(2)$ -invariant hermitian structure.
3. If the anti-self-dual metric is specified by a solution of Painlevé V with one real and two complex parameters, or Painlevé II with one real parameter, then for any $x \in M$ there exists one real null surface passing through x .
4. If the anti-self-dual metric is specified by a solution of Painlevé III with two real parameters, then for any $x \in M$ there exist two real null surfaces passing through x .

7 Appendix

We review the Painlevé equations, second order nonlinear differential equations without moving critical points. We list six equations classified by Painlevé and Gambier, where $\alpha, \beta, \gamma, \delta$ are parameters [9].

1. Painlevé I

$$\frac{d^2 q}{dx^2} = 6q^2 + x.$$

2. Painlevé II

$$\frac{d^2 q}{dx^2} = 2q^3 + xq + \alpha.$$

3. Painlevé III

$$\frac{d^2 q}{dx^2} = \frac{1}{q} \left(\frac{dq}{dx} \right)^2 - \frac{1}{x} \frac{dq}{dx} + \frac{1}{x} (\alpha q^2 + \beta) + \gamma q^3 + \frac{\delta}{q}.$$

4. Painlevé IV

$$\frac{d^2 q}{dx^2} = \frac{1}{2q} \left(\frac{dq}{dx} \right)^2 + \frac{3}{2} q^3 + 4xq^2 + 2(x^2 - \alpha)q + \frac{\beta}{q}.$$

5. Painlevé V

$$\frac{d^2 q}{dx^2} = \left(\frac{1}{2q} + \frac{1}{q-1} \right) \left(\frac{dq}{dx} \right)^2 - \frac{1}{x} \frac{dq}{dx} + \frac{(q-1)^2}{x^2} \left(\alpha q + \frac{\beta}{q} \right) + \frac{\gamma q}{x} + \frac{\delta q(q+1)}{q-1}.$$

6. Painlevé VI

$$\begin{aligned} \frac{d^2 q}{dx^2} = & \frac{1}{2} \left(\frac{1}{q} + \frac{1}{q-1} + \frac{1}{q-x} \right) \left(\frac{dq}{dx} \right)^2 - \left(\frac{1}{x} + \frac{1}{x-1} + \frac{1}{q-x} \right) \frac{dq}{dx} \\ & + \frac{q(q-1)(q-x)}{x^2(x-1)^2} \left\{ \alpha + \beta \frac{x}{q^2} + \gamma \frac{x-1}{(q-1)^2} + \delta \frac{x(x-1)}{(q-x)^2} \right\}. \end{aligned}$$

References

- [1] Atiyah, M. F. and Hitchin, N. J.: Low energy scattering of non-Abelian monopoles, *Phys. Lett. A* **107** (1985), 21–25
- [2] Atiyah, M. F., Hitchin, N. J. and Singer, I. M.: Self-duality in four-dimensional Riemannian geometry, *Proc. Roy. Soc, London Ser. A* **362** (1978) 425–461
- [3] Belinski, V. A., Gibbons, G. W., Page, D. W., Page, D. W. and Pope, C. N.: Asymptotically Euclidean Bianchi IX metrics in quantum gravity, *Phys. Lett. B* **76** (1978), 433–435
- [4] Dancer, A. S.: Scalar-flat Kähler metrics with $SU(2)$ symmetry, *J. reine. angew. Math.* **479** (1996), 99–120
- [5] Jimbo, M., Miwa, T. and Ueno, K.: Monodromy Preserving Deformation of Linear Ordinary Differential Equations with Rational Coefficients. I, *Physica* **2D** (1981), 306–352
- [6] Jimbo, M. and Miwa, T.: Monodromy Preserving Deformation of Linear Ordinary Differential Equations with Rational Coefficients. II, *Physica* **2D** (1981), 407–448
- [7] Hitchin, N. J.: Twistor spaces, Einstein metrics and isomonodromic deformations, *J. Differential Geom.* **42** (1995), 30–112
- [8] Okumura, S.: The Self-Dual Hermitian Metric and Painlevé III, preprint
- [9] Painlevé, P.: Sur les équations différentielles du second ordre à points critiques fixes, *C. R. Acad. Sci. Paris* **143** (1906), 1111–1117
- [10] Penrose, R.: Nonlinear gravitons and curved twistor theory, *Gen. Rel. Grav.* **7** (1976), 31–52
- [11] Tod, K. P.: Self-dual Einstein metrics from the Painlevé VI equation, *Phys. Lett. A* **190** (1994), 221–224